

PROCEEDINGS OF THE TWENTY-SIXTH ANNUAL KECK RESEARCH SYMPOSIUM IN GEOLOGY

April 2013
Pomona College, Claremont, CA

Dr. Robert J. Varga, Editor
Director, Keck Geology Consortium
Pomona College

Dr. Jade Star Lackey
Symposium Convener
Pomona College

Carol Morgan
Keck Geology Consortium Administrative Assistant

Christina Kelly
Symposium Proceedings Layout & Design
Office of Communication & Marketing
Scripps College

*Keck Geology Consortium
Geology Department, Pomona College
185 E. 6th St., Claremont, CA 91711
(909) 607-0651, keckgeology@pomona.edu, keckgeology.org*

ISSN# 1528-7491

The Consortium Colleges

The National Science Foundation

ExxonMobil Corporation

**KECK GEOLOGY CONSORTIUM
PROCEEDINGS OF THE TWENTY-SIXTH ANNUAL KECK RESEARCH
SYMPOSIUM IN GEOLOGY**

ISSN# 1528-7491

April 2013

Robert J. Varga
Editor and Keck Director
Pomona College

Keck Geology Consortium
Pomona College
185 E 6th St., Claremont, CA
91711

Christina Kelly
Proceedings Layout & Design
Scripps College

Keck Geology Consortium Member Institutions:

**Amherst College, Beloit College, Carleton College, Colgate University, The College of Wooster,
The Colorado College, Franklin & Marshall College, Macalester College, Mt Holyoke College,
Oberlin College, Pomona College, Smith College, Trinity University, Union College,
Washington & Lee University, Wesleyan University, Whitman College, Williams College**

2012-2013 PROJECTS

TECTONIC EVOLUTION OF THE CHUGACH-PRINCE WILLIAM TERRANE: SHUMAGIN ISLANDS AND KENAI PENINSULA, ALASKA

Faculty: *JOHN GARVER*, Union College, *CAMERON DAVIDSON*, Carleton College

Students: *MICHAEL DELUCA*, Union College, *NICOLAS ROBERTS*, Carleton College, *ROSE PETTIETTE*, Washington & Lee University, *ALEXANDER SHORT*, University of Minnesota-Morris, *CARLY ROE*, Lawrence University.

LAVAS AND INTERBEDS OF THE POWDER RIVER VOLCANIC FIELD, NORTHEASTERN OREGON

Faculty: *NICHOLAS BADER & KIRSTEN NICOLAYSEN*, Whitman College.

Students: *REBECCA RODD*, University of California-Davis, *RICARDO LOPEZ-MALDONADO*, University of Idaho, *JOHNNY RAY HINOJOSA*, Williams College, *ANNA MUDD*, The College of Wooster, *LUKE FERGUSON*, Pomona College, *MICHAEL BAEZ*, California State University-Fullerton.

BIOGEOCHEMICAL CARBON CYCLING IN FLUVIAL SYSTEMS FROM BIVALVE SHELL GEOCHEMISTRY - USING THE MODERN TO UNDERSTAND THE PAST

Faculty: *DAVID GILLIKIN*, Union College, *DAVID GOODWIN*, Denison University.

Students: *ROXANNE BANKER*, Denison University, *MAX DAVIDSON*, Union College, *GARY LINKEVICH*, Vassar College, *HANNAH SMITH*, Rensselaer Polytechnic Institute, *NICOLLETTE BUCKLE*, Oberlin College, *SCOTT EVANS*, State University of New York-Geneseo.

METASOMATISM AND THE TECTONICS OF SANTA CATALINA ISLAND: TESTING NEW AND OLD MODELS

Faculty: *ZEB PAGE*, Oberlin College, *EMILY WALSH*, Cornell College.

Students: *MICHAEL BARTHELMES*, Cornell College, *WILLIAM TOWBIN*, Oberlin College, *ABIGAIL SEYMOUR*, Colorado College, *MITCHELL AWALT*, Macalester College, *FREDY, AGUIRRE*, Franklin & Marshall College, *LAUREN MAGLIOZZI*, Smith College.

GEOLOGY, PALEOECOLOGY AND PALEOCLIMATE OF THE PALEOGENE CHICKALOON FORMATION, MATANUSKA VALLEY, ALASKA

Faculty: *CHRIS WILLIAMS*, Franklin & Marshall College, *DAVID SUNDERLIN*, Lafayette College.

Students: *MOLLY REYNOLDS*, Franklin & Marshall College, *JACLYN WHITE*, Lafayette College, *LORELEI CURTIN*, Pomona College, *TYLER SCHUETZ*, Carleton College, *BRENNAN O'CONNELL*, Colorado College, *SHAWN MOORE*, Smith College.

CRETACEOUS TO MIOCENE EVOLUTION OF THE NORTHERN SNAKE RANGE METAMORPHIC CORE COMPLEX: ASSESSING THE SLIP HISTORY OF THE SNAKE RANGE DECOLLEMENT AND SPATIAL VARIATIONS IN THE TIMING OF FOOTWALL DEFORMATION, METAMORPHISM, AND EXHUMATION

Faculty: *MARTIN WONG*, Colgate University, *PHIL GANS*, University of California-Santa Barbara.

Students: *EVAN MONROE*, University of California-Santa Barbara, *CASEY PORTELA*, Colgate University, *JOSEPH WILCH*, The College of Wooster, *JORY LERBACK*, Franklin & Marshall College, *WILLIAM BENDER*, Whitman College, *JORDAN ELMIGER*, Virginia Polytechnic Institute and State University.

THE ROLE OF GROUNDWATER IN THE FLOODING HISTORY OF CLEAR LAKE, WISCONSIN

Faculty: *SUSAN SWANSON*, Beloit College, *JUSTIN DODD*, Northern Illinois University.

Students: *NICHOLAS ICKS*, Northern Illinois University, *GRACE GRAHAM*, Beloit College, *NOA KARR*, Mt. Holyoke College, *CAROLINE LABRIOLA*, Colgate University, *BARRY CHEW*, California State University-San Bernardino, *LEIGH HONOROF*, Mt. Holyoke College.

PALEOENVIRONMENTAL RECORDS AND EARLY DIAGENESIS OF MARL LAKE SEDIMENTS: A CASE STUDY FROM LOUGH CARRA, WESTERN IRELAND

Faculty: *ANNA MARTINI*, Amherst College, *TIM KU*, Wesleyan University.

Students: *SARAH SHACKLETON*, Wesleyan University, *LAURA HAYNES*, Pomona College, *ALYSSA DONOVAN*, Amherst College.

INTERDISCIPLINARY STUDIES IN THE CRITICAL ZONE, BOULDER CREEK CATCHMENT, FRONT RANGE, COLORADO

Faculty: David Dethier, Williams College, Will Ouimet, U. Connecticut.

Students: *CLAUDIA CORONA*, Williams College, *HANNAH MONDRACH*, University of Connecticut, *ANNETTE PATTON*, Whitman College, *BENJAMIN PURINTON*, Wesleyan University, *TIMOTHY BOATENG*, Amherst College, *CHRISTOPHER HALCSIK*, Beloit College.

Funding Provided by:
Keck Geology Consortium Member Institutions
The National Science Foundation Grant NSF-REU 1062720
ExxonMobil Corporation

**Keck Geology Consortium: Projects 2012-2013
Short Contributions— Matanuska, Alaska Project**

GEOLOGY, PALEOECOLOGY AND PALEOCLIMATE OF THE PALEOGENE CHICKALOON FORMATION, MATANUSKA VALLEY, ALASKA

Faculty: *CHRIS WILLIAMS*, Franklin & Marshall College, *DAVID SUNDERLIN*, Lafayette College.

CLIMATE AND PLANT INTERACTIONS IN AN ANCIENT ALASKAN RAINFOREST

MOLLY REYNOLDS, Franklin and Marshall College

Research Advisor: Christopher Williams

GROWTH RING ANALYSIS OF *METASEQUOIA* FROM THE NORTHEAST USA AND THE PALEOCENE/EOCENE CHICKALOON FORMATION, ALASKA

JACLYN WHITE, Lafayette College

Research Advisor: David Sunderlin

ZIRCON U-PB GEOCHRONOLOGY FROM THE PALEOCENE-EOCENE CHICKALOON FORMATION

LORELEI CURTIN, Pomona College

Research Advisor: Robert Gaines

EXAMINING THE PALEOENVIRONMENT OF THE CHICKALOON FORMATION, MATANUSKA VALLEY, ALASKA THROUGH *CAMPELOMA* AND *UNIONOIDA* FOSSIL ASSEMBLAGES

TYLER SCHUETZ, Carleton College

Research Advisor: Clint Cowan

USING STABLE ISOTOPE GEOCHEMISTRY TO RECONSTRUCT THE PALEOHYDROLOGY AND PALEOCLIMATE OF THE PALEOCENE-EOCENE CHICKALOON FORMATION, ALASKA

BRENNAN O'CONNELL, Colorado College

Research Advisor: Henry Fricke

PROVENANCE ANALYSIS OF MEDIAL BASIN CHANNEL SANDS IN THE CHICKALOON FORMATION, MATANUSKA VALLEY, ALASKA

SHAWN A. MOORE, Smith College, Northampton MA

Research Advisor: Bosiljka Glumac

EXAMINING THE PALEOENVIRONMENT OF THE CHICKALOON FORMATION, MATANUSKA VALLEY, ALASKA THROUGH *CAMPELOMA* AND *UNIONOIDA* FOSSIL ASSEMBLAGES

TYLER SCHUETZ, Carleton College

Research Advisor: Clint Cowan

INTRODUCTION

The Chickaloon Formation is located in the Matanuska Valley Region of Alaska, approximately 70 miles northeast of Anchorage. The Chickaloon is a >1500m thick terrestrial sedimentary unit and is composed of a complex succession of sandstones, laminated siltstones, mudstones, carbonaceous shales, coal beds and conglomerates (Neff et al., 2011; Sunderlin et al., 2011). The purpose of this research was to determine more about higher latitude continental settings preserved in this unit during and near the Paleocene-Eocene thermal maximum. The formation has been dated using ash layers that approximately straddle the Paleocene-Eocene boundary and previous research has determined that the formation represents sedimentation in a high-latitude (~60°) temperate environment (Sunderlin et al., 2011). In addition to abundant plant remains, the Chickaloon Formation contains a variety of gastropod and bivalve fossils, which further constrain the paleoenvironment of the formation to be a meandering floodplain setting (Neff et al., 2011). This claim is supported by the modern analogues of *Campeloma* snails and *Unionoida* bivalves collected and analyzed in this project

METHODS

In my examination of the invertebrate fauna of the Chickaloon Formation I mapped a horizontally continuous exposure of the unit at the Evan Jones Mine and collected samples of both gastropod and bivalve fossils. Fossiliferous beds were located using previous knowledge of the area and stratigraphic sections were measured surrounding the beds for approximately half a meter. This project focused on one primary mollusk fossil bed.

All specimens were brought back to Carleton College and were labeled and measured, with particular attention given to the spacing and orientation of visible growth lines on the fossil surface. The specimens were identified through a review of the literature and the observed associations between certain gastropod and bivalve species.

I then studied modern analogues of the identified fossils and the mapped stratigraphic sections of the formation to constrain the depositional environment of the Chickaloon Formation at the Evan Jones Mine.

RESULTS

Approximately 50 samples were catalogued and identified, and were tied to three distinct locations along the sedimentological/environmental gradient. These samples were identified using morphological measurements and comparison with paleoecological literature and with modern analogues. Particular attention was paid to observed modern associations between gastropods and bivalves. The gastropods were identified as members of the Viviparidae family, and the genus *Campeloma*. The bivalves were identified as members of the *Unionoida* family and the Unionoidae family. This identification, as well as the stratigraphic analysis of the stratigraphic columns supports the claim that the paleoenvironment of the Chickaloon Formation is represented by a floodplain, fluvial setting with periodic crevasse splays resulting from episodic flood events.

DISCUSSION

Facies Associations

Two main facies were observed in the stratigraphic columns taken at the Evan Jones Mine location. The first layer is dominated by organic material, with heavily fractured clay interbeds and carbonaceous shale and mudstone; the top of the section is marked by grey to tan sandstone beds. In general the sections coarsen upward, which is indicative of crevasse splays in a fluvial environment. Figures 1, 2, and 3 show the observed stratigraphy at each location.




Figure 1. Stratigraphic column of West vertical section. This column is comprised of interbedded coal and clay layers coarsening upward to poorly consolidated sandstones.




Figure 2. Stratigraphic column of center vertical section. This column contains an ash layer, which was not visible at other locations. The sand layers are much more lithified than in other columns.




Figure 3. Stratigraphic column of East vertical section. This section shows much larger organic layers and thinner sandstone layers.

Facies Association I – organic layers**Description**

3.5cm of coal, in contact with 2 cm of a fine grained, heavily fractured clay. The coal layer continues above the clay for 9 cm, and becomes less consolidated and siltier toward the top of the layer. Occasional presence of an ash layer, but this is not continuous across all measured columns. As the beds move eastward, they become much less lithified.

Interpretation

The presence of interbedded layers of coal and shale is indicative of a mire environment, where organic material is preserved under anoxic conditions. The variations of shale beds to coal beds occur when the mire floods, which causes an influx of material into the system, forming shale beds.

Facies Association II – coarsening upward sandstone**Description**

Grey, carbonaceous shale and mudstone continue for 11 cm and are in contact with grey sandstone. The sandstone is very fine grained, with bivalve and gastropod fossils found in the contact between carbonaceous shale and sand layers. Above the grey sandstone lies 4 cm of tan sandstone that contains iron rich concretions. This sandstone is less fine grained and the interactions between the two continue upward out of the study section. The exposure in general coarsens upward, however the sandstone beds fine upward. Moving to the east along the exposure, the sandstones begin to form grey and orange sandstone ribbons. All specimens collected were found in the contact between facies I and II.

Interpretations

This succession was deposited in slow moving waters in a lacustrine or fluvial environment. Crevasse splays are represented by the sandstone beds that laterally pinch and swell, and were caused by sand deposited in overbank deposits during flood conditions. The increasing grain size represents the increase in carrying capacity of the fluvial environment the stratigraphic column continued I would expect to see an abrupt termination of increasing grain size indicating when the flood conditions concluded.

Taxonomy/Modern Analogs**Modern Genus *Viviparus***

Modern *Viviparus*, as described in Richardson and Brown (1989) are detritivores who populate shallow waters with pH ranges of 6.3-8.5 (Duch, 1976; Vincent, 1979; Jokinen and Pondick, 1981). *Viviparus* typically lives for 18-48 months (Eckblad and Shealy, 1972; Richardson and Brown, 1989). *Viviparus* is often common in areas that experience non-energetic flow and not much current but have sufficient water movement to maintain adequate oxygen concentration for the gastropods to survive (Richardson and Brown, 1989). Often found in eutrophic lakes, ponds, and slow-moving streams, *Viviparus* prefers to silt-mud areas (Duch, 1979). Hanley and Flores (1984) found that *Viviparus* prefers large, shallow, low energy environment with low clastic influx.

Fossil Genus *Campeloma*

Twenty-five *Campeloma* fossil remains were identified by their low to moderate spire, impressed sutures, and dextral nature (Jokinen, 1992; Thorp, 2001). The shell is thick, with a rounded or convex center spiral and a simple, continuous peristome. The surface of the shell is smooth, and the only striae are growth lines (Ellsworth, 1886). The species *Campeloma* is generally found in fluvial environments and is common in lowland river courses, often with bivalves like *Unio*, *Anodonta*, *Margaritifera*, and *Coricula* (Yen, 1951).

Campeloma are most often found at the muddy bottoms of ponds, rivers, and creeks, are intolerant of temperatures higher than 90 degrees, and are sensitive to changes in mineral character of the water (Ellsworth, 1886).

Campeloma is a genus of the Viviparidae family, whose taxonomy has been discussed more than *Campeloma*. Therefore, to constrain my inferred paleoenvironment, I will assume *Campeloma* is analogous to modern day *Viviparus* and that their environmental preferences are similar.

Campeloma specimens were found exclusively in Facies Association II – the coarsening upward sandstone at the study area. This facies is comprised of different sandstone layers that pinch and swell

laterally, indicative of a crevasse splay deposition. *Campeloma* favors large, shallow, low energy environments and could have been entrapped when the crevasse splay spilled out onto the low energy flood plain. Crevasse-splay deposits occur when large quantities of floodwater and sediment are diverted into adjacent flood basins by following distinct channels cut across the banks through natural levee deposits (Reineck and Singh, 1973). These channels are called crevasses, and when water overtops the levees of the main channel crevasse splay systems can develop their own distributary channel pattern (Reineck and Singh, 1973).

Ellsworth (1887) reports finding frost-killed specimens, indicating that *Campeloma* responds drastically to changes in environment. *Campeloma* is used as a bioremediation indicator because it reacts to changes sediment composition, implying it must live in relatively stable environments. This conclusion is supported by the observed disappearance of *Campeloma* communities in areas that receive sewage from cities or manufactories. The paleoenvironment of the *Campeloma* fossils must have been stable during the life span of the gastropod; temperatures must not have exceeded 90° and the pH range must have been constrained to 6.3-8.5.

Order Unionoida

I interpret the Chickaloon bivalve specimens to be members of the Order Unionoida based on a comparison of the literature (Vaughn and Hakenkamp, 2001, Thorp and Covich, 2001, and Vaughn and Spooner, 2006). Unionids typically inhabit stable, freshwater habitats (Thorp and Covich, 2001). Mussel beds tend to be prominent in areas that are stable during periods of high water inundation, or flooding (Vaughn and Spooner, 2006). *Unionoida* are found in patchy distributions, caused by patchily distributed resources (Vaughn and Spooner, 2006). Vaughn and Hakenkamp (2001) observed the impact Unionoida have on their environments and found that the bivalves are integral to maintaining certain environmental conditions.

At my study area, unionids are found in the contact between the organic layers and sandstones. This indicates Unionoida survives in a variety of stable, freshwater environments. Unionoidae specimens were

found in the crevasse splay deposition, along with *Campeloma*. This association has been recognized in the literature before, and is supported by the observation that mussel beds are more common in areas of flooding or inundation.

Growth Line Analysis

The abundance and pattern of growth lines can help constrain seasonality and variation in paleoenvironment. Growth lines are used to determine the age of a specimen at death. However, the thickness of growth lines depends on the climatic factors during the period of growth (Imbrie and Newell, 1964). In many skeletonized organisms, growth lines grow by accretion over one or more years. Seasonal environment variations influence this growth, so we can cautiously interpret growth lines as annual in nature (Ivany, 2012). Butler et al. (2010) argue that growth lines in mollusk and gastropod fossil are annual and can be used to construct chronology for marine environments that are analogous to those produced from tree rings in the terrestrial environment. Growth line variations potentially record paleoclimate indicators such as temperature, salinity, nutrients and dissolved oxygen (Carter, 1980). Therefore, examining growth lines of molluscan and gastropod specimens are a potential way to constrain the paleoclimate of an area.

Growth line patterns of the observed specimens (both *Campeloma* and Unionoids) show periodicity in groupings. The lines are arranged into two distinct sets, one with many closely spaced thin lines, and one with cm scale spacing between lines. These patterns indicate seasonality in the paleoenvironment, with closely spaced line indicating shorter growing seasons with less available water and greater spacing showing longer seasons with more available water (possibly flood like conditions). These patterns indicate the differences in seasons were consistent. The lack of growth lines on some areas of the specimens and the raised varices indicate halts in growth, which could be due to changes in seasonality and climate. More research should be done into the possibility of further constraining seasonality with growth line analysis. Unfortunately, the specimens collected in this project did not have original preserved material, so the analysis could not be completed.

CONCLUSIONS

By studying the distributions, tolerances, and environments of modern day *Campeloma* and *Unionoidae* we can start to recreate the paleoenvironment of the specimens found at Evan Jones Mine. Both *Campeloma* and *Unionoidae* are found in stable, fresh-water, fluvial environments. *Campeloma* prefer large, shallow low energy environments and *Unionoidae* are most prominent in areas with stable conditions analogous to flood plains. These preferences along with observed stratigraphy indicate the Chickaloon Formation experienced periodic episodes of crevasse splay deposition. By looking at the modern and fossilized examples of *Campeloma* and *Unionoidae* I sought to determine if the fossil assemblages I observed in the field matched the previous paleoclimate research on the Chickaloon Formation. I found that the Chickaloon was deposited in a fluvial, crevasse plain with evidence for the presence of mires and floodplains. These observations correlated well with previous research on the area, supporting the view of the Chickaloon as a high-latitude continental fluvial setting. This research contributes to the available knowledge of the paleoclimate during the Paleocene-Eocene thermal maximum by lending more credibility to already published theories. Further research in this area should focus on the observed periodicity of growth intervals to further constrain patterns of temperature and seasonality.

REFERENCES CITED:

- Butler, P.G., Richardson, C.A., Scourse, J.D., Wanamaker, D.J., Shammon, T.M., and Bennell, J.D., 2010, Marine climate in the Irish Sea: analysis of a 489-year marine master chronology derived from growth increments in the shell of the clam *Arctica islandica*. *Quaternary Science Reviews*, 29:1614-1632.
- Carter, J.G., 1980, Environmental and biological controls of bivalve shell mineralogy and microstructure: New York, Plenum Press, 113 p.
- Duch, T.M., 1976, Aspects of the feeding habits of *Viviparus georgianus*: *The Nautilus*, 90:7-10.
- Eckblad, J.W., and Shealy, M.H., 1972, Predation on largemouth bass embryos by the pond snail *Viviparus georgianus*, *Transactions of the American Fisheries Society*, 101:734-738.
- Ellsworth, R., 1886., On the genus *Campeloma*, Rafinesque, with a revision of the species, recent and fossil, *Bulletin of the Washburn College Laboratory of Natural History*, 1:149-174.
- Hanley, J.H., and Flores, R.M., 1987, Taphonomy and paleoecology of nonmarine Mollusca: Indicators of alluvial plain lacustrine sedimentation, upper part of the Tongue River member, Fort Union Formation (Paleocene), northern Powder River basin, Wyoming and Montana, *PALAIOS*, 2:479-496.
- Imbrie, J., and Newell, N., 1964, Approaches to paleoecology: New York, John Wiley & Sons, Inc., 432 p.
- Ivany, L.C., 2012, Reconstructing paleoseasonality from accretionary skeletal carbonates – challenges and opportunities, *The Paleontological Society Papers*, 18:133-165.
- Jokinen, E.H., and Pondick, J., 1981, Rare and endangered species: Freshwater gastropods of southern New England, *The Bulletin of the American Malacological Union*, 50:52-53.
- Leeder, M., 2011, Sedimentology and sedimentary basins: from turbulence to tectonics: New Jersey, Wiley-Blackwell, 768 p.
- Neff, J.L., Hagadorn, J.W., Sunderlin, D., and Williams, C.J., 2011, Sedimentology, facies architecture and chemostratigraphy of a continental high-latitude Paleocene-Eocene succession – The Chickaloon Formation, Alaska, *Sedimentary Geology*, 240:14-29.
- Reineck, H.E., and Singh, I.B., 1973, Depositional sedimentary environments: New York, Springer-Verlag, 439 p.
- Richardson, T.D., and Brown, K.M., 1989, Secondary production of two subtropical snails (*prosobranchia: Viviparidae*), *Journal of the North American Benthological Society*, 8:229-236.
- Sunderlin, D., Loope, G., Parker, N.E., Williams, C.J., 2011, Paleoclimate and paleoecology implications of a Paleocene-Eocene fossil leaf assemblage, Chickaloon Formation, Alaska. *PALAIOS*, 26:335-345.
- Thorp, J.H., and Covich, A.P., 2001, Ecology and classification of North American freshwater invertebrates: Academic Press, 1056 p.
- Vaughn, C.C., and Hakenkamp, C.C., 2001, The func-

- tional role of burrowing bivalves in freshwater ecosystems: *Freshwater Biology*, 46:1431-1446.
- Vaughn, C.C, and Spooner, D.E., 2006, Unionid mussels influence macro-invertebrate assemblage structure in streams: *Journal of the North American Benthological Society*, 25:691-700.
- Vincent, B., 1979, Etude du benthos d'eau douce dans le haut-estuaire de Saint-Laurent (Quebec): *Canadian Journal of Zoology*, 57: 1271-2182.
- Yen, T.C., 1951, Fossil fresh-water mollusks and ecological interpretations, *Bulletin of the Geological Society of America*, 62:1375-1380.